

$g/u(1)^d$ parafermions from constrained WZNW theories.

A.V.Bratchikov *

Kuban State Technological University,
2 Moskovskaya Street, Krasnodar, 350072, Russia.

December, 1997

Abstract

The conformal field theory based on the $g/u(1)^d$ coset construction is treated as the WZNW theory for the affine Lie algebra $\hat{g}$ with the constrained $\hat{u}(1)^d$ subalgebra. Using a modification of the generalized canonical quantization method generators and primary fields of an extended symmetry algebra are found for arbitrary d .

1 Introduction.

The $g/u(1)^d$, $1 \leq d \leq \text{rank } g$, parafermions are generators of an extended symmetry algebra of the conformal field theory based on the $g/u(1)^d$ coset construction [1, 2]

$$K = L^g - L^{u(1)^d}. \quad (1)$$

where L^g is the energy-momentum tensor, which is associated with the affine Lie algebra $\hat{g}$ via the Sugawara formula. In the case of $su(2)/u(1)$ the parafermions were constructed by Fateev and Zamolodchikov [3]. These authors found equations which relate the $su(2)/u(1)$ parafermions with the $\hat{su}(2)$ algebra. A generalization of these equations to all simple Lie algebra was

*bratchikov@elect.ru

proposed in ref.[4].However it was not proved that the generalized equations define parafermions.

In a recent article [5] we displayed the approach to the study of the $g/u(1)^d$ conformal field theory which is based on the connection between this theory and the Wess-Zumino-Novikov-Witten (WZNW) theory for $\hat{g}$ with the constrained $\hat{u}(1)^d$ currents.It was shown using an operator version of Dirac's procedure [6] that the initial energy-momentum tensor L^g is replaced in the constrained theory by K . It turned out that the constrained $su(2)$ WZNW theory is equivalent to the $su(2)/u(1)$ theory.However in the general case the constrained currents determine the $g/u(1)^d$ parafermions only up to some factors.

In the present paper we choose another way. We perform operator quantization of the constrained WZNW theory using the approach which is based on a modification of the generalized canonical quantization method (see e.g.[7] and references therein) and show that the resulting symmetry algebra is an extended symmetry algebra of the $g/u(1)^d$ coset conformal field theory.We present two explicit realizations for the parafermions: a realization where they are represented as elements of a coset algebra and a vertex operator construction.Using the second realization we find the equations which define the $g/u(1)^d$ parafermions through their relationship with the generators of $\hat{g}$.The theory has a discrete symmetry group which is generated by the $u(1)^r$ currents, where $r = rank g$.The quantization method enables us to find primary fields of the $g/u(1)^d$ theory.A representation of the parafermionic algebra is obtained by applying the parafermions to the parafermionic primary fields.

The parafermionic algebra is associated with a classical quadratically non-linear algebra which can be obtained using both the generalized canonical quantization method and the Dirac-bracket formalism.This demonstrates the equivalence of the quantization methods.

The plan of the paper as follows.In Sec.2 we describe the quantization method.Putting auxiliary operators into correspondence with first and second class constraints we convert the initial constraints into effective abelian constraints. The constrained algebra is obtained by replacing the initial operators by operators which commute with the effective abelian constraints.In section 3 we review the WZNW theory and introduce the constraints.In Sec.4 we perform quantization of the constrained WZNW theory and find operators which replace the initial symmetry generators. These operators are

expressed in terms of the initial currents and free bosonic fields. We show that the $g/u(1)^d$ parafermionic algebra can be formulated as a coset algebra in a natural way and find the parafermionic currents and primary fields. In Sec.6 we find vertex operator construction for the $g/u(1)^d$ parafermionic algebra. In Sec.7 we consider the constrained classical current algebra, find the corresponding Dirac brackets and show that the same brackets can be obtained using the generalized canonical quantization method.

2 The quantization method.

Let T^A and T_μ be first and second class constraints. For our purpose it is sufficient to consider constraints of zero Grassman parity $\epsilon(T^A) = \epsilon(T_\mu) = 0$ and assume that they obey the commutator relations

$$[T^A, T^B] = [T^A, T_\mu] = 0, \quad [T_\mu, T_\nu] = r_{\mu\nu}, \quad (2)$$

where $r_{\mu\nu}$ is an invertible c-number matrix.

According to the prescription of the generalized canonical quantization method [7] we introduce the auxiliary operators b_μ which obey the commutator relation

$$[b_\mu, b_\nu] = r_{\mu\nu} \quad (3)$$

and convert the second class constraints T_μ into the effective abelian constraints

$$\tilde{T}_\mu = T_\mu + ib_\mu. \quad (4)$$

It is easy to see that

$$[\tilde{T}_\mu, \tilde{T}_\nu] = 0, \quad \tilde{T}_\mu(0) = T_\mu \quad (5)$$

and

$$[\tilde{T}_\mu, T^A] = 0. \quad (6)$$

Let us put into correspondence with each constraint T^A a pair of the canonical operators (p^A, q^A) which, in contrast with ref. [7], are supposed to be of the same Grassman parity as the constraints. In the extended phase space the operator T^A can be replaced by an operator $\tilde{T}^A(\chi)$, $\chi = (b_\mu, p^A, q^A)$, which, as well as T^A , satisfies the equations

$$[\tilde{T}_\mu, \tilde{T}^A] = 0, \quad [\tilde{T}^A, \tilde{T}^B] = 0 \quad (7)$$

and the boundary conditions

$$\tilde{T}^A(0) = T^A. \quad (8)$$

Let F be an initial operator. We shall demand that F is replaced in the constrained theory by an operator $\mathcal{F}(\chi)$ which commutes with the effective abelian constraints $(\tilde{T}^A, \tilde{T}_\mu)$. The generating equations of the constrained algebra then assume the form

$$\begin{aligned} [\tilde{T}_\mu, \mathcal{F}] &= 0, & [\tilde{T}^A, \mathcal{F}] &= 0, \\ \mathcal{F}(0) &= F. \end{aligned} \quad (9)$$

Solutions of equations (7) and (9) are not completely fixed by the boundary conditions. The arbitrariness is to be removed by a gauge condition.

3 The WZNW model and constraints.

The WZNW theory [8, 9] is invariant with respect to two commuting affine Lie algebras, which are generated by the left and right currents $(H^s(z), E^\alpha(z))$, $(\bar{H}^s(\bar{z}), \bar{E}^\alpha(\bar{z}))$, where $s = 1, \dots, r$ and α are roots of g . Since the algebras generated by the left and right currents are identical we shall only consider the first. The theory is also invariant with respect to the conformal algebra. The Virasoro generator $L^g(z)$ is expressed quadratically in terms of the currents.

The fields $H^s(z)$, $E^\alpha(z)$ and $L^g(z)$ satisfy the operator product expansions

$$\begin{aligned} H^s(z)H^t(w) &= \frac{k\delta^{st}}{(z-w)^2}, & H^s(z)E^\alpha(w) &= \frac{\alpha^s E^\alpha(w)}{z-w}, \\ E^\alpha(z)E^\beta(w) &= \begin{cases} \frac{f(\alpha, \beta)E^{\alpha+\beta}(w)}{z-w}, & \text{if } \alpha + \beta \text{ is a root} \\ \frac{k}{(z-w)^2} + \frac{\alpha \cdot H(w)}{z-w}, & \text{if } \alpha + \beta = 0 \\ 0, & \text{otherwise} \end{cases} \\ L^g(z)H^s(w) &= \frac{H^s(w)}{(z-w)^2} + \frac{\partial_w H^s(w)}{z-w}, \\ L^g(z)E^\alpha(w) &= \frac{E^\alpha(w)}{(z-w)^2} + \frac{\partial_w E^\alpha(w)}{z-w}, \\ L^g(z)L^g(w) &= \frac{c_g}{2(z-w)^4} + \frac{2L^g(w)}{(z-w)^2} + \frac{\partial_w L^g(w)}{z-w}, \end{aligned}$$

$$c_g = \frac{k \dim g}{k + h^\vee}, \quad (10)$$

where k is the level of the representation, $f(\alpha, \beta)$ are the structure constants and $h^\vee$ is the dual Coxeter number of the algebra g .

Let g be simply-laced. In this case the currents $H^s(z)$, $E^\alpha(z)$ and energy-momentum tensor $L^g(z)$ can be expressed in terms of the bosonic fields

$$\varphi^{sj}(z) = x^{sj} - i a_o^{sj} \log z + i \sum_{n \neq 0} \frac{a_n^{sj}}{n} z^{-n}, \quad (11)$$

where $j = 1, \dots, k$, and

$$[x^{sj}, a_o^{tl}] = i \delta^{st} \delta^{jl}, \quad [a_m^{sj}, a_n^{tl}] = m \delta^{st} \delta^{jl} \delta_{m+n, 0}. \quad (12)$$

The bosonic construction for $H^s(z)$, $E^\alpha(z)$ and $L^g(z)$ read [10, 11, 12]

$$\begin{aligned} E^\alpha(z) &= \sum_{j=1}^k : e^{i\alpha \cdot \varphi^j(z)} : c_\alpha^j, & H_s(z) &= \sum_{j=1}^k i \partial_z \varphi^{sj}(z), \\ L^g(z) &= \frac{1}{2(k + h^\vee)} \left((1 + h^\vee) \sum_{s=1}^r \sum_{j=1}^k : (i \partial_z \varphi^{sj})^2 : \right. \\ &\quad \left. + 2 \sum_{s=1}^r \sum_{i < j}^k : (i \partial_z \varphi^{sj})(i \partial_z \varphi^{si}) : + 2 \sum_{\alpha} \sum_{i < j}^k : \exp(i\alpha \cdot (\varphi^i - \varphi^j)) : c_\alpha^i c_{-\alpha}^j \right), \end{aligned} \quad (13)$$

where $::$ denotes normal ordering with respect to the modes of the bosons, $\alpha^2 = 2$ and c_α^j is a cocycle operator.

We shall consider the WZNW theory subject to the constraints

$$H^A(z) \approx 0, \quad (14)$$

where $A = 1, \dots, d$. It is convenient to decompose the constraints into modes

$$\begin{aligned} H^A(z) &= H_o^A z^{-1} + \sum_{n \neq 0} H_n^A z^{-n-1}, \\ H_o^A &= \sum_{j=1}^k a_o^{Aj}, & H_n^A &= \sum_{j=1}^k a_n^{Aj}, \end{aligned} \quad (15)$$

and consider an equivalent set of constraints:

$$H_o^A \approx 0, \quad H_n^A \approx 0. \quad (16)$$

The operators H_o^A, H_n^A obey the algebra

$$[H_o^A, H_o^B] = [H_o^A, H_n^B] = 0, \quad [H_m^A, H_n^B] = km\delta^{AB}\delta_{m+n,0}. \quad (17)$$

It follows from these commutator relations that the constraints H_o^A are first class and H_n^A are second class.

4 Quantization of the model.

4.1 Generating equations of the constrained symmetry algebra.

To quantize the system we put into correspondence with the constraints the operators $\chi = (b_n^A, p^A, q^A)$ which satisfy the commutator relations

$$[b_m^A, b_n^B] = km\delta^{AB}\delta_{m+n,0}, \quad [q^A, p^B] = ik\delta^{AB}. \quad (18)$$

The effective abelian constraints $\tilde{H}_n^A(b)$ are given by

$$\tilde{H}_n^A = H_n^A + ib_n^A \quad (19)$$

and we replace the first class constraints H_o^A by the following

$$\tilde{H}_o^A = H_o^A - ip^A. \quad (20)$$

According to eq.(9) the other operators of the constrained symmetry algebra $\mathcal{H}^P(z) \equiv \mathcal{H}^P(\chi, z)$, where $P = d+1, \dots, r$, $\mathcal{E}^\alpha(z) \equiv \mathcal{E}^\alpha(\chi, z)$ and $\mathcal{L}^g(z) \equiv \mathcal{L}^g(\chi, z)$ satisfy the equations

$$[\tilde{H}_N^A, \mathcal{H}^P] = 0, \quad [\tilde{H}_N^A, \mathcal{E}^\alpha] = 0, \quad [\tilde{H}_N^A, \mathcal{L}^g] = 0 \quad (21)$$

with the boundary conditions

$$\mathcal{H}^P(0, z) = H^P(z), \quad \mathcal{E}^\alpha(0, z) = E^\alpha(z), \quad \mathcal{L}^g(0, z) = L^g(z). \quad (22)$$

A solution of these equations is given by

$$\begin{aligned}\mathcal{H}^P(z) &= H^P(z), & \mathcal{E}^\alpha(z) &= E^\alpha(z) : e^{\frac{1}{k}\tilde{\alpha}\cdot\phi(z)} :, \\ \mathcal{L}^g(z) &= K(z) + \frac{1}{2k} \left(\tilde{H}^A(z) \right)^2,\end{aligned}\tag{23}$$

where $\tilde{\alpha} = (\alpha^A)$, K is given by (1),

$$\tilde{H}^A(z) = H^A(z) + \partial_z \phi^A(z)\tag{24}$$

and $\phi = (\phi^A)$ are the bosonic fields:

$$\phi^A(z) = q^A - ip^A \log z + i \sum_{n \neq 0} \frac{b_n^A}{n} z^{-n}.\tag{25}$$

This solution is not the general one. It can be fixed by appropriate gauge conditions.

4.2 Parafermionic currents.

The operator product expansions of K with $\mathcal{H}^P$ and $\mathcal{E}^\alpha$ are given by

$$\begin{aligned}K(z)\mathcal{H}^P(w) &= \frac{\mathcal{H}^P(w)}{(z-w)^2} + \frac{\partial_w \mathcal{H}^P(w)}{z-w}, \\ K(z)\mathcal{E}^\alpha(w) &= \frac{\Delta_\alpha \mathcal{E}^\alpha(w)}{(z-w)^2} + \frac{1}{z-w} \left(\partial_w \mathcal{E}^\alpha(w) - \frac{1}{k} \tilde{\alpha} \cdot \tilde{H}(w) \mathcal{E}^\alpha(w) \right),\end{aligned}\tag{26}$$

where

$$\Delta_\alpha = 1 - \frac{\tilde{\alpha}^2}{2k}\tag{27}$$

Let Ω be the algebra generated by $\mathcal{H}^P(z)$, $\mathcal{E}^\alpha(z)$ and $K(z)$. Since these operators commute with $\tilde{H}_N^A$ so therefore must all the fields of Ω . Let us define the following set of the fields

$$\Upsilon = (U \in \Omega; U|_{\tilde{H}^A=0} = 0).\tag{28}$$

One can check that for arbitrary $U(z) \in \Upsilon$ and $X(z) \in \Omega$

$$U(z)X(w) \in \Upsilon.\tag{29}$$

Hence Υ is an invariant subalgebra of Ω and the coset space Ω/Υ is an algebra. We shall denote by $\{X\}(z)$ the coset represented by the field $X(z)$.

The coset field $\{K\}(z)$ satisfies the $g/u(1)^d$ Virasoro algebra:

$$\{K\}(z)\{K\}(w) = \frac{c_{g/u(1)^d}}{2(z-w)^4} + \frac{2\{K\}(w)}{(z-w)^2} + \frac{\partial_w\{K\}(w)}{z-w}, \quad (30)$$

where $c_{g/u(1)^d} = c_g - d$. It follows from (23) that $\{K\}(z)$ can be represented by $\mathcal{L}^g(z)$ as well. Equations (26) express the fact that $\{\mathcal{H}^P\}(z)$ and $\{\mathcal{E}^\alpha\}(z)$ are primary fields of the $g/u(1)^d$ Virasoro algebra:

$$\begin{aligned} \{K\}(z)\{\mathcal{H}^P\}(w) &= \frac{\{\mathcal{H}^P\}(w)}{(z-w)^2} + \frac{\partial_w\{\mathcal{H}^P\}(w)}{z-w}, \\ \{K\}(z)\{\mathcal{E}^\alpha\}(w) &= \frac{\Delta_\alpha\{\mathcal{E}^\alpha\}(w)}{(z-w)^2} + \frac{\partial_w\{\mathcal{E}^\alpha\}(w)}{z-w}, \end{aligned} \quad (31)$$

The theory has the discrete symmetry group which is generated by H_o^s :

$$[H_o^s, \{\mathcal{H}^P\}] = 0, \quad [H_o^s, \{\mathcal{E}^\alpha\}] = \alpha^s\{\mathcal{E}^\alpha\} \quad [H_o^s, \{K\}] = 0. \quad (32)$$

Let us consider the case of $su(2)/u(1)$. The $\hat{su}(2)$ generators $E^+(z), E^-(z)$ and $H(z)$ are given by

$$\begin{aligned} E^+(z) &= \sum_{j=1}^k : e^{i\sqrt{2}\varphi^j(z)} :, & E^-(z) &= \sum_{j=1}^k : e^{-i\sqrt{2}\varphi^j(z)} :, \\ H(z) &= \sum_{j=1}^k i\partial_z\varphi^j(z), \end{aligned} \quad (33)$$

where $\varphi^j \equiv \varphi^{1j}$. In the theory with the constrained current $H(z) \approx 0$ these operators are replaced by

$$\begin{aligned} \mathcal{E}^+(z) &= E^+(z) : e^{\frac{\sqrt{2}}{k}\phi(z)} :, & \mathcal{E}^-(z) &= E^-(z) : e^{-\frac{\sqrt{2}}{k}\phi(z)} :, \\ \tilde{H}(z) &= H(z) + \partial_z\phi(z). \end{aligned} \quad (34)$$

We find that

$$\begin{aligned} \mathcal{E}^+(z)\mathcal{E}^-(w) &= k(z-w)^{-2+\frac{2}{k}} \left(I + \tilde{h}(w)(z-w) \right. \\ &\quad \left. + \left(\frac{k+2}{k}K(w) + \frac{1}{2}\partial_w\tilde{h}(w) + \frac{1}{2}\tilde{h}^2(w) \right)(z-w)^2 \right), \end{aligned} \quad (35)$$

where $\tilde{h} = \tilde{H}/k$ and K is given by the $su(2)/u(1)$ coset construction. From this it follows that the coset currents $\{\mathcal{E}^+\}(z)$ and $\{\mathcal{E}^-\}(z)$ satisfy the parafermionic algebra of ref.[3]

$$\{\mathcal{E}^+\}(z)\{\mathcal{E}^-\}(w) = k(z-w)^{-2+\frac{2}{k}} \left(\{I\} + \frac{k+2}{k} \{K\}(w)(z-w)^2 \right). \quad (36)$$

4.3 Primary fields.

Let $G_\lambda^\Lambda(z)$ be the initial primary field with highest weight Λ which has the weights λ

$$\begin{aligned} E_n^\alpha G_\lambda^\Lambda &= H_n^s G_\lambda^\Lambda = L_n^g G_\lambda^\Lambda = 0, & n > 0, \\ H_o^s G_\lambda^\Lambda &= \lambda^s G_\lambda^\Lambda, & E_o^\alpha G_\lambda^\Lambda = t_{\lambda\lambda'}^\alpha G_{\lambda'}^\Lambda, \\ L_o^g G_\lambda^\Lambda &= \Delta_\Lambda G_\lambda^\Lambda, & L_{-1}^g G_\lambda^\Lambda = \partial_z G_\lambda^\Lambda, \end{aligned} \quad (37)$$

where t^α are the step generators of g in the representation with highest weight Λ . The anomalous dimension Δ_Λ is given by [9]

$$\Delta_\Lambda = \frac{\Lambda(\Lambda + \rho)}{2(k + h^\vee)}, \quad (38)$$

where ρ is half the sum positive roots of g .

Let us define

$$\mathcal{G}_\lambda^\Lambda(z) = G_\lambda^\Lambda(z) : e^{\frac{1}{k} \tilde{\lambda} \cdot \phi(z)} :, \quad (39)$$

where $\tilde{\lambda} = (\lambda^A)$. This field satisfies the following equations

$$\begin{aligned} \tilde{H}_n^A \mathcal{G}_\lambda^\Lambda &= 0, & n \geq 0, & \quad K_n \mathcal{G}_\lambda^\Lambda = 0, & n > 0, \\ K_o \mathcal{G}_\lambda^\Lambda &= \Delta_\lambda^\Lambda \mathcal{G}_\lambda^\Lambda, & K_{-1} \mathcal{G}_\lambda^\Lambda &= \partial_z \mathcal{G}_\lambda^\Lambda - \frac{1}{k} \tilde{\lambda} \cdot \tilde{H}_{-1} \mathcal{G}_\lambda^\Lambda, \end{aligned} \quad (40)$$

where

$$\Delta_\lambda^\Lambda = \Delta_\Lambda - \frac{\tilde{\lambda}^2}{2k}. \quad (41)$$

Let T be the space which is obtained by applying the currents of Ω repeatedly to the fields $\mathcal{G}_\lambda^\Lambda$. Let

$$V = (\mathcal{G} \in T; \mathcal{G}|_{\tilde{H}^A=0} = 0). \quad (42)$$

The space V is an invariant subspace with respect to the algebra Ω . Hence one can consider the representation of Ω in the coset space T/V . It follows from eq.(40) that the coset $\{\mathcal{G}_\lambda^\Lambda\}$ which is represented by $\mathcal{G}_\lambda^\Lambda$ is the primary field with the anomalous dimension Δ_λ^Λ (41).

The space T/V can be decomposed according to the transformation properties under the $u(1)^r$ algebra. The field $\{\mathcal{G}_\lambda^\Lambda\}$ belongs to the subspace with the charge λ :

$$H_o^s\{\mathcal{G}_\lambda^\Lambda\} = \lambda^s\{\mathcal{G}_\lambda^\Lambda\}. \quad (43)$$

5 Vertex operator construction for the parafermionic algebra.

In this section we show that the theory for g simply-laced can be formulated in terms of the initial operators x^{sj} , a_o^{sj} , and a_n^{sj} .

The constraints (19) and (20) can be considered as the strong operator equations

$$p^A = \sum_{j=1}^k a_o^{Aj} \quad b_n^A = \sum_{j=1}^k a_n^{Aj}. \quad (44)$$

Substituting this into (23) we express the constrained currents in terms of the initial operators and the auxiliary operators $q = (q^A)$

$$\mathcal{E}^\alpha(z) = \mathcal{E}_o^\alpha(z) e^{\frac{1}{k}\tilde{\alpha}\cdot q}, \quad (45)$$

where

$$\mathcal{E}_o^\alpha(z) = \sum_{j=1}^k : e^{i[\tilde{\alpha}\cdot\omega^j(z) + \tilde{\alpha}^\perp\cdot\varphi^j(z)]} : c_\alpha^j, \quad (46)$$

$$\omega^{Aj}(z) = x^{Aj} - i\eta^{jl} \left(a_o^{Al} \log z - \sum_{n \neq 0} \frac{a_n^{Al}}{n} z^{-n} \right) \quad (47)$$

and

$$\eta^{ij} = \begin{cases} \frac{(k-1)}{k} & \text{if } i = j, \\ -\frac{1}{k} & \text{if } i \neq j. \end{cases} \quad (48)$$

Note that $\mathcal{E}^\alpha(z)$ (45) belongs to the coset $\{\mathcal{E}^\alpha\}(z)$.

Let us put into correspondence with $\{K\}, \{\mathcal{H}^P\}$ and $\{\mathcal{E}^\alpha\}$ the fields K, H^P and $\mathcal{E}^\alpha$ (45) respectively. This map is the isomorphism of the algebras. Since q commute with all the initial operators we can set $q = 0$ in (45). Therefore the generators of Ω/Υ can be represented by the fields K, H^P and $\mathcal{E}_o^\alpha$ which are expressed in terms of x^{sj}, a_o^{sj} , and a_n^{sj} . We have checked that $\mathcal{E}_o^\alpha(z)$ satisfies the equation

$$K(z)\mathcal{E}_o^\alpha(w) = \frac{\Delta_\alpha \mathcal{E}_o^\alpha(w)}{(z-w)^2} + \frac{\partial_w \mathcal{E}_o^\alpha(w)}{z-w}. \quad (49)$$

We find that the initial currents $E^\alpha(z)$ can be expressed as follows

$$E^\alpha(z) =: e^{\frac{i}{k}\tilde{\alpha}\cdot\tilde{\varphi}(z)} : r^\alpha \mathcal{E}_o^\alpha(z), \quad (50)$$

where

$$r^\alpha = e^{-\frac{i}{k}\tilde{\alpha}\cdot x}, \quad x^A = \sum_{j=1}^k x^{Aj} \quad (51)$$

and

$$\tilde{\varphi}^A = \sum_{j=1}^k \varphi^{Aj} \quad (52)$$

These equations can be used as the generating equations for the $g/u(1)^d$ parafermions.

In the case of $su(2)$ the cocycles c_α^j can be replaced by the identity operator. Since x^1 commutes with $\mathcal{E}_o^\alpha(z)$ we can set $x^1 = 0$ and rewrite equations (50) in the form [3]

$$\begin{aligned} E^+(z) &= \sqrt{k}\psi^-(z) : e^{i\frac{\sqrt{2}}{k}\tilde{\varphi}(z)} :, \\ E^-(z) &= \sqrt{k}\psi^+(z) : e^{-i\frac{\sqrt{2}}{k}\tilde{\varphi}(z)} :, \end{aligned} \quad (53)$$

where

$$\psi^-(z) = \frac{1}{\sqrt{k}}\mathcal{E}_o^+(z) \quad \psi(z) = \frac{1}{\sqrt{k}}\mathcal{E}_o^-(z) \quad (54)$$

and $\tilde{\varphi}(z) \equiv \tilde{\varphi}^1(z)$.

These results can be generalized to non-simply-laced algebras using the vertex operator representation of the associated affine Lie algebras [13, 14].

6 Classical parafermions and the Dirac-bracket formalism.

Quantization of the $g/u(1)^d$ theory can also be performed by using the Dirac-bracket formalism. The approach presented above is equivalent to the canonical quantization of the classical WZNW theory.

To show this we first find a classical system which corresponds to the $g/u(1)^d$ theory. Let us replace the operators $H^s(z)$, $E^\alpha(z)$ and $\phi^A(z)$ by the functions $H^s(\sigma)$, $E^\alpha(\sigma)$, $\phi^A(\sigma)$, satisfying the Poisson bracket relations

$$\begin{aligned} \{H^s(\sigma), H^t(\sigma')\} &= -k\delta^{st}\delta', & \{H^s(\sigma), E^\alpha(\sigma')\} &= \alpha^s E^\alpha(\sigma)\delta, \\ \{E^\alpha(\sigma), E^\beta(\sigma')\} &= \begin{cases} f(\alpha, \beta)E^{\alpha+\beta}(\sigma)\delta, & \text{if } \alpha + \beta \text{ is a root} \\ -k\delta' + \alpha \cdot H(\sigma)\delta, & \text{if } \alpha + \beta = 0 \\ 0, & \text{otherwise} \end{cases}, \end{aligned} \quad (55)$$

$$\{\phi^A(\sigma), \phi^B(\sigma')\} = \frac{k}{2}\delta^{AB}\epsilon, \quad (56)$$

where $\delta = \delta(\sigma - \sigma')$ and $\epsilon = \epsilon(\sigma - \sigma')$. From this it follows that the constraints $H^A(\sigma) \approx 0$ are second class. The abelian operator constraints $\tilde{H}^A(z)$ are replaced by $\tilde{H}^A(\sigma) = H^A(\sigma) + \partial_\sigma \phi(\sigma)$ which satisfy the relations

$$\{\tilde{H}^A(\sigma), \tilde{H}^B(\sigma')\} = 0. \quad (57)$$

It is easy to check that the functions

$$\mathcal{H}^P(\sigma) = H^P(\sigma), \quad \mathcal{E}^\alpha(\sigma) = E^\alpha(\sigma)e^{\frac{1}{k}\tilde{\alpha} \cdot \phi(\sigma)} \quad (58)$$

commute with $\tilde{H}^A(\sigma)$

$$\{\tilde{H}^A(\sigma), \mathcal{H}^P(\sigma')\} = 0, \quad \{\tilde{H}^A(\sigma), \mathcal{E}^\alpha(\sigma')\} = 0. \quad (59)$$

Using the Poisson bracket relations (55) and (56) we find

$$\begin{aligned} \{\mathcal{H}^P(\sigma), \mathcal{H}^Q(\sigma')\} &= -k\delta^{PQ}\delta', & \{\mathcal{H}^P(\sigma), \mathcal{E}^\alpha(\sigma')\} &= \alpha^P \mathcal{E}^\alpha(\sigma)\delta, \\ \{\mathcal{E}^\alpha(\sigma), \mathcal{E}^\beta(\sigma')\} &= \begin{cases} f(\alpha, \beta)\mathcal{E}^{\alpha+\beta}(\sigma)\delta + \\ + \frac{\tilde{\alpha} \cdot \tilde{\beta}}{2k}\mathcal{E}^\alpha(\sigma)\mathcal{E}^\beta(\sigma')\epsilon, & \text{if } \alpha + \beta \text{ is a root} \\ -k\delta' + \tilde{\alpha} \cdot \tilde{H}(\sigma)\delta + \tilde{\alpha}^\perp \cdot \mathcal{H}(\sigma)\delta - \\ - \frac{\tilde{\alpha}^2}{2k}\mathcal{E}^\alpha(\sigma)\mathcal{E}^{-\alpha}(\sigma')\epsilon, & \text{if } \alpha + \beta = 0 \\ \frac{\tilde{\alpha} \cdot \tilde{\beta}}{2k}\mathcal{E}^\alpha(\sigma)\mathcal{E}^\beta(\sigma')\epsilon, & \text{otherwise,} \end{cases} \end{aligned} \quad (60)$$

where $\mathcal{H}(\sigma) = (\mathcal{H}^P(\sigma))$. Similar algebra was found by the authors of ref.[15] using the gauged WZNW lagrangian.

The algebra R defined by equations (57),(59) and (60) can be obtained using the Dirac-bracket formalism. Let I and J belong to the initial current algebra (55). In the constrained theory they are replaced by the currents which satisfy the Dirac bracket relations

$$\{I, J\}_D = \{I, J\} - \int d\sigma d\sigma' \{I, H^A(\sigma)\} \{H^A(\sigma), H^B(\sigma')\}^{-1} \{H^B(\sigma'), J\}. \quad (61)$$

We find that H^s and E^α obey the algebra

$$\begin{aligned} \{H^A(\sigma), H^B(\sigma')\}_D &= \{H^A(\sigma), H^P(\sigma')\}_D = \{H^A(\sigma), E^\alpha(\sigma')\}_D = 0, \\ \{H^P(\sigma), H^Q(\sigma')\}_D &= -k\delta^{PQ}\delta', \quad \{H^P(\sigma), E^\alpha(\sigma')\}_D = \alpha^P E^\alpha(\sigma)\delta, \\ \{E^\alpha(\sigma), E^\beta(\sigma')\}_D &= \begin{cases} f(\alpha, \beta) E^{\alpha+\beta}(\sigma)\delta + \\ + \frac{\tilde{\alpha} \cdot \tilde{\beta}}{2k} E^\alpha(\sigma) E^\beta(\sigma') \epsilon, & \text{if } \alpha + \beta \text{ is a root} \\ -k\delta' + \alpha \cdot H(\sigma)\delta - \\ - \frac{\tilde{\alpha}^2}{2k} E^\alpha(\sigma) E^{-\alpha}(\sigma') \epsilon, & \text{if } \alpha + \beta = 0 \\ \frac{\tilde{\alpha} \cdot \tilde{\beta}}{2k} E^\alpha(\sigma) E^\beta(\sigma') \epsilon, & \text{otherwise} \end{cases} \end{aligned} \quad (62)$$

This algebra is isomorphic to R . The corresponding map is

$$H^A(\sigma) \rightarrow \tilde{H}^A(\sigma), \quad H^P(\sigma) \rightarrow \mathcal{H}^P(\sigma), \quad E^\alpha(\sigma) \rightarrow \mathcal{E}^\alpha(\sigma). \quad (63)$$

To quantize the system we could therefore start from the Dirac brackets (62) (or equivalently from the Poisson brackets (57),(59) and (60)), use ansatz (58) and replace the functions $H^s(\sigma)$, $E^\alpha(\sigma)$ and $\phi^A(\sigma)$ by the corresponding operators.

7 Conclusion.

Treating the $g/u(1)^d$ conformal field theory as the WZNW theory for $\hat{g}$ with the constrained $\hat{u}(1)^d$ currents and using a modification of the generalized canonical quantization method we have found the explicit construction of the symmetry generators of the $g/u(1)^d$ theory. The presented quantization may be generalized to the WZNW theory for $\hat{g}$ with an arbitrary constrained current algebra $\hat{h} \subset \hat{g}$. We expect that in the general case the conformal invariant solution of the generating equations determines a symmetry algebra of the g/h conformal field theory.

References

- [1] K.Bardakci and M.B.Halpern,*Phys.Rev.* **D3**(1971)2493.
M.B.Halpern,*Phys.Rev.* **D4**(1971)2398.
- [2] P.Goddard,A.Kent and D.Olive,*Phys.Lett.* **152B** (1985) 88.
- [3] V.A.Fateev and A.B.Zamolodchikov,*Sov.Phys.JETP* **82** (1985) 215.
- [4] D.Gepner,*Nucl.Phys.* **B290** (1987) 10.
- [5] A.V.Bratchikov,*Phys.Lett.***420B**(1998) 285, hep-th/9710036.
- [6] P.A.M.Dirac,*Lectures in Quantum Mechanics* (Yeshiva University, New York,1964).
- [7] I.A.Batalin,E.S.Fradkin and T.Fradkina,*Nucl.Phys.* **B332** (1990) 723.
- [8] E.Witten,*Comm.Math.Phys.* **92** (1984) 455.
- [9] V.Knizhnik and A.B.Zamolodchikov,*Nucl.Phys.* **B247** (1984) 83.
- [10] I.B.Frenkel and V.G.Kac,*Inv.Math.***62**(1980)23.
- [11] G.Segal,*Commun.Math.Phys.* **80** (1981) 301.
- [12] G.V.Dunne,I.G.Halliday and P.Suranyi,*Nucl.Phys.* **B325** (1989) 526.
- [13] P.Goddard,W.Nahm,D.Olive and A.Schwimmer, *Commun.Math.Phys.* **107** (1986) 179.
- [14] D.Bernard and J.Thierry-Mieg, *Commun.Math.Phys.* **111** (1987) 181.
- [15] K.Bardakci,M.Crescimanno and E.Rabinovici,*Nucl.Phys.* **B344** (1990) 344.